Identification Notes for wildlife law enforcement

BI	RI	
	BI	BIRI

Scientific Name/s: Falconiformes

Common Name/s: eagle, falcon, hawk, vulture

PROBLEM: Mounted or whole specimens of raptors are presented for identification.

Diurnal birds of prey (Falconiformes) are recognized by their hooked beaks and robust feet. Well-developed talons are commonly present. Determination to Family is the first step in the identification process. When country or region of origin is known, most diurnal raptors can be assigned to genus (and often to species) by reference to field guides. The following key will assist you in assigning fledgling and adult diurnal birds of prey to Family.

IT WILL NOT WORK FORHATCHLINGS AND CHICKS.










no bony tubercle no tomial tooth

(in aside view of a perforate nostril, you can see through the bird's bill)

KEY TO FAMILIES OF DIURNAL BIRDS OF PREY

a. tarsus shorter than 8 inches


b, tarsus longer than 8 inches Sagittariidae: Secretary Bird (Africa)



2. a. nostrils **imperforate**, hind toe not elevated, hind **talon** well developed.......3







b. nostrils perforate, hind toe elevated and reduced........... Cathartidae: New World vultures (North, South and Central America)







New World vulture

PLEASE TURN OVER

Dentification Notes

CATEGORY:

BIRD

Common Name


Falconiformes eagle, falcon, hawk, vulture



bony tubercle


tomial tooth









b. no **tomial** tooth on bill, no bony **tubercle** in nostril Accipitridae: kites, hawks, eagles and Old World vultures (worldwide except Antarctica)











Accipiter



Old World vulture

REFERENCES:

Newton, I., ed. 1990. Birds of Prey. Facts on File, Inc. New York. 240 pp. Weick, F. 1980. Birds of Prey of the World. Collins, London. 159 pp.

Illustrations by: Anthony E. Kerwin